

Distal Biceps Tendon Repair Rehabilitation Guidelines

Name	: Date:
Diagn	osis: Date of Surgery:
□ im	MOBILIZATION Posterior splint Elbow immobilization at 90 degrees for 7-10 days
BF	RACE Elbow placed in ROM brace at 7-10 days post-operative ROM set at 45 degrees to full flexion Gradually increase elbow ROM in brace
	Brace or sling is discontinued at 8 weeks . ANGE OF MOTION PROGRESSION
•	Week 2: 45 degrees to full flexion Week 3: 45 degrees to full flexion Week 4: 30 degrees to full flexion Week 5: 20 degrees to full flexion Week 6: 10 degrees to full flexion; full supination/pronation Week 8: full ROM of elbow; full supination/pronation
RA	ANGE OF MOTION EXERCISES
•	Week 3-5: passive ROM for elbow flexion and supination; active-assisted ROM for elbow extension and pronation Week 6-8: initiate AAROM elbow flexion Week 9: active ROM elbow flexion
Ce	enter for Musculoskeletal Care • 333 Fast 38h Street, New York, NY 1001 • 7646\ 501-7070, • cmc med nyu edu



STRENGTHENING PROGRAM

- Week 2-3: isometrics for triceps and shoulder muscles
- Week 9: active ROM, no resistance applied
- Week 10: PRE (progressive resistive exercise) program is initiated for elbow flexion and supination/pronation
 - o Begin with one pound and gradually increase shoulder strengthening program
- Week 14: may initiate light weight training such as bench press and shoulder press

Comments:		
Frequency:times per week	Duration:weeks	
Signature:		Date